

FIG. 1-1

Constitutively Active Receptors

File Name	Receptor	Mutation Site	Sequence	Assay / Cells	Reference
CLASS A GROUP I					
MSHR_mouse	melanocyte-stimulating hormone	TMII	92 VSIVLETTIIL K	adenyl cyclase activity/ HEK293, stably transfected	(Robbins, Nadeau et al. 1993)
MSH					
CLASS A GROUP II SH1B_human	5-hydroxytryptamine _{1B}	C-terminus of IC3	313 RERKATKTLGI K, R, Q	binding of [³⁵ S]GTP[S] / CHO-K1	(Paywels, Gouble et al. 1999)
SH2A_human	5-hydroxytryptamine _{2A}	C-terminus of IC3	322 NEQKAKVLGI K	IP production / COS-7	(Egan, Herrick-Davis et al. 1998)
2H2C_rat	5-hydroxytryptamine _{2C}	C-terminus of IC3	312 NEDDAKVLGI L	PI hydrolysis / COS-7	(Herrick-Davis, Egan et al. 1997)

FIG. 1-2

CLASS A GROUP II					
A1AB_human	α_{1B} -adrenergic alpha 1B-AR	TMDI junction between TMDIII and IC2	63 A FAIVGNILVIL SEQ ID NO: 6 142 A CAISIDRYIGV SEQ ID NO: 7	IP / COS-7	(Scheer, Fanelli et al. 1997)
A1AB_human	α_{1B} -adrenergic alpha 1B-AR	junction between TMDIII and IC2	143 K CAISIDRYIGV SEQ ID NO: 8	IP / COS-7	(Scheer, Costa et al. 2000)
A1AB_human	α_{1B} -adrenergic	TMIII carboxyl end of IC3 TMV	128 F AVDVLCTASI SEQ ID NO: 9 293 E REKKA A KT L GI SEQ ID NO: 10 204 V EPPFYALFSSLG SEQ ID NO: 11	IP / COS-1 IP arachidonic acid release IP / COS-1	(Perez, Hwa et al. 1996) (Hwa, Gaivin et al. 1997)
A1AB_human	α_{1B} -adrenergic	C-terminal IC3	293 SREKKA A KT SEQ ID NO: 12 X=19 different substitutions	PI / COS-7	(Kjelsberg, Cotecchia et al. 1992)
A1AB_human	α_{1B} -adrenergic	C-terminus IC3	288 293 K H L KFSREKKA A KT L GI SEQ ID NO: 13	PI hydrolysis / rat fibroblast	(Allen, Lefkowitz et al. 1991)
A2AA_human	α_2 C10-adrenergic alpha-2AAR	C-terminal IC3 loop	373 (348?) EKRF T VLAV SEQ ID NO: 14 X=F, A, C, E, K	adenyl cyclase inhibition / HEK293	(Ren, Kurose et al. 1993)
ACM1_human	muscarinic Hm1 muscarinic acetylcholine M1	C-terminal IC3 loop junction	360 A SLVKEKKA A RTLS SEQ ID NO: 15	PI / HEK(U293)	(Högger, Shockey et al. 1995)
ACM2-human	muscarinic acetylcholine M2	junction of IC3 and TMVI	390 KKVTRTIL 1 A SEQ ID NO: 16 1-4 A inserted	IP production, inhibition of cAMP production / COS-7	(Liu, Blin et al. 1996)

FIG. 1-3

CLASS A GROUP II			TMVI				
ACM3_rat	m3 muscarinic (rat) muscarinic acetylcholine M3		TMVI			507 TWTPYNIMVLVNT SEQ ID NO: 17 S	IP / COS-7 (Bliiml, Muischler et al. 1994)
ACM5_human	m5 muscarinic muscarinic acetylcholine M5		N-terminus to TMII TMVI			chimera composed of m2 1-69 m5 77-445 m2 391-466	(Burststein, Spalding et al. 1996)
ACM5_human	m5 muscarinic muscarinic acetylcholine M5		TMVI	SEQ ID NO: 18		451 459 465 AILLA EITW TPVNI MVLVST M L H C V S F T	(Spalding, Burststein et al. 1998)
ACM5_human	m5 muscarinic muscarinic acetylcholine M5		junction of TMVI and EC3			465 YNIMVLVSTFCDKCV SEQ ID NO: 19 X=V,F,R,K,+more	(Spalding, Burststein et al. 1997)
B1AR_human	β_1 -adrenergic		C-terminus			389 RKAFOGLLCCA SEQ ID NO: 20 R	(Mason, Moore et al. 1999)
D2AR_human	β_2 -adrenergic beta-2AR		C-terminal IC3 loop			266 272 FCLKEHKALKTLGI SEQ ID NO: 21 SR K A	(Samama, Cotecchia et al. 1993); (Lefkowitz, Cotecchia et al. 1993)
DADR_human	dopamine DIA		carboxyl terminal IC3			264 SFKMSEKRETKVLKT SEQ ID NO: 22 I K 288 from D1B receptor APDTSIKKETKVLKT SEQ ID NO: 23	(Charpentier, Jarvie et al. 1996)
DADR_human	dopamine DI		TMVI			286 FVCCWLPFFIL SEQ ID NO: 24 A	(Cho, Taylor et al. 1996)
HH2R_rat	histamine H ₂		IC2			115 FMISLDRYCAV SEQ ID NO: 25 N, A	(Alewijns, Timmerman et al. 2000)

4/31

FIG. 1-4

File Name	Receptor	Mutation Site	Sequence	Assay / Cells	Reference
CLASS A GROUP III					
OPSD_human	opsin	TMII	⁹⁰ FMVLGGFTSTLY SEQ ID NO: 26 D	transducin; phosphorylation by rhodopsin kinase / COS	(Rim and Oprian 1995)
	rhodopsin	TMIII	¹¹³ GCNLEGGFFAT SEQ ID NO: 27 Q		
		TMVII	^{292 296} MTIPAFFAKSAAY SEQ ID NO: 28 E G, E, M ²⁹³ Ala neutral a.a converted to carboxylate and competes with ¹¹³ Glu for salt bridge with ²⁹⁶ Lys		
OPSD_human	opsin	TMIII	¹³⁴ VVLAIERYVWV SEQ ID NO: 29 I, Q, S	transducin; radioligand binding / COS	(Acharya and Kamik 1996)
OPSD_human	rhodopsin				
OPSD_human	opsin	TM6	²⁵⁷ RMVIIMVIAFL SEQ ID NO: 30 Y, N	transducin, GTPγS uptake / COS	(Han, Smith et al. 1998)
	rhodopsin	plus TM3	plus G113Q		
OPSD_human	opsin	TMVII	²⁹⁶ PAFFAKSAAY SEQ ID NO: 31 G X=E,M natural mutants + 10 different a.a. substitutions	transducin; radioligand binding / COS	(Govardhan and Oprian 1994); (Cohen, Yang et al. 1993)
	rhodopsin		disrupts critical salt bridge between ²⁹⁶ Lys(TMVII) and ¹¹³ Glu(TMIII)		
		IC2	¹³⁴ VVLAIERYVWV SEQ ID NO: 32 Q		(Cohen, Yang et al. 1993)

5/31




FIG. 1-5

TRFR_mouse	thyrotropin-releasing hormone TRH-R	carboxyl tail	335 FRKLGNCKQK STOP	SEQ ID NO: 33	⁴⁵ Ca ²⁺ efflux, [Ca ²⁺] / Xenopus oocytes; IP formation / A1T20, <i>stably transfected</i>	(Matus-Leibovitch, Nussenzeig et al. 1995)
------------	--	---------------	---------------------------	---------------	--	---

FIG. 1-6

6/31

File Name	Receptor	Mutation Site	Sequence	Assay / Cells	Reference
CLASS A GROUP IV					
BRB2_human	bradykinin B ₂ B2 bradykinin BK-2	TMIII TMVI	¹¹³ ATISMNLYSSI SEQ ID NO: 34 A ²⁵⁶ LLPFICWLPFQI SEQ ID NO: 35 F	IP production / COS-7	(Marie, Koch et al. 1999)

2010501 5496M007



FIG. 1-7

File Name	Receptor	Mutation Site	Sequence	Assay / Cells	Reference
CLASS A GROUP V					
AG2R_rat	AT _{1A} Type-1A angiotensin II	TMIII	111 ASVSENLVASV SEQ ID NO: 36 A disrupts ¹¹¹ Asn(TMIII) - ¹¹² Tyr(TMVII) interaction	phospholipase C; IP production / COS-7	(Grobowski, Maigret et al. 1997)
AG2R_rat	AT _{1A}	C-terminus of TM7	305 LFYGF LGKKFK SEQ ID NO: 37 Q	IP production / HEK-293; intracellular Ca ²⁺ mobilization / CHO	(Parnot, Bardin et al. 2000)
FMRLR_human	Type-1A angiotensin II formylmethionylleucylphenylalanine (fMLPR)	IC1 other multiple mutations	51 LVIVVAGFRMTHTVTTISYNKAVA LVVVVTAFAEAKRTINAIWFLNLAVA (K above conflicts with SWISS-PROT database)	PI production; phospholipase C stimulation / COS-7	(Amatruda, Dragas-Graonic et al. 1995)
IL8B_human	interleukin-8 receptor B CXCR-2 chemokine	IC2	138 ACISVD RYLAIVH SEQ ID NO: 40 V	IP production; Ca ²⁺ mobilization and actin polymerization / NIH 3T3	(Burger, Burger et al. 1999)
LSHR_human	luteinizing hormone (LH)	IC3	564 MATNKDTKI AKK SEQ ID NO: 41 G	cAMP production / HEK293	(Kudo, Osuga et al. 1996)
LSHR_human	luteinizing hormone (LH)	TMVI	578 ILIFTD FTCMA SEQ ID NO: 42 G	cAMP production / COS-7	(Shenker, Laue et al. 1993)
LSHR_human	luteinizing hormone (LH)	TM6 SEQ ID NO: 43	571 577 KI AKKMAILLIFTDFTCM I I	cAMP production / COS-7	(Kosugi, Van Dop et al. 1995)
LSHR_rat	luteinizing hormone / human chorionic gonadotropin (LH/hCG)	TMVI	556 ILIFTD FTCMA SEQ ID NO: 44 G, Y	cAMP production / HEK 293T	(Bradbury, Kawate et al. 1997; Bradbury and Menon 1999)
OPRD_mouse	delta opioid receptor	TM3	128 KVLSD YNNMF SEQ ID NO: 45 A, K, H	adenylyl cyclase inhibition / COS-7	(Cavalli, Babey et al. 1999)
OXYR_human	oxytocin	IC2	137 LMSLD RCLAIC SEQ ID NO: 46 A	IP production / COS-7	(Fanelli, Barbier et al. 1999)

FIG. 1-8

PAFR_human	platelet-activating factor (PAF)	C-terminus of IC3	231 EVKRRALWVCTVLAV SEQ ID NO: 47 R	IP production / COS-7	(Parent, Le Gouill et al. 1996)
PAFR_human	platelet-activating factor (PAF)	TMIII	100 CLFFINTYCSV SEQ ID NO: 48 A	arachidonate release, IP production, adenyllyl cyclase inhibition / CHO	(Ishii, Izumi et al. 1997)
PE23_human	prostaglandin E ₂ , EP3III EP3IV	C-terminal tail	360 FCQEEFWGN SEQ ID NO: 49 FCQMRKRRLREOEEFWGN SEQ ID NO: 50 ↑truncated	inhibition of adenyllyl cyclase / CHO-K1	(Jin, Mao et al. 1997)
PE23_mouse	prostaglandin E ₂ , EP3	carboxyl-terminal tail SEQ ID NO: 51	336 KILLRKFCQIRDHT (3α) MMNHL (3β) ↑truncated	inhibition of adenylate cyclase / CHO, stably expressed	(Hasegawa, Negishi et al. 1996)
THR _R _human	thrombin	EC2 loop SEQ ID NO: 52	259 CHDVLNETLEGGYVY DLKD KDF I 268	⁴⁵ Ca ²⁺ efflux, PI hydrolysis, reporter gene induction / COS-7	(Nanevicz, Wang et al. 1996)
TSHR_human	thyrotropin (TSHR) thyroid stimulating hormone	EC1	486 YYNHAIDWQTG SEQ ID NO: 53 F, M	inositol phosphate-- diacylglycerol cascade / COS-7	(Parma, Van Sande et al. 1995)
		EC2	568 YAKVSI CLPMD SEQ ID NO: 54 T		
TSHR_human	thyrotropin (TSHR) thyroid stimulating hormone	TMIII	509 ASELSVYTLTV SEQ ID NO: 55 A	adenyllyl cyclase activation / COS-7	(Duprez, Parma et al. 1994)
		TMVII	672 YPLNSCANPFL SEQ ID NO: 56 Y		
TSHR_human	thyrotropin (TSHR) thyroid stimulating hormone	TMV	597 VAFVIVCCCHV SEQ ID NO: 57 L	cAMP formation / COS-7 cells	(Esapa, Duprez et al. 1999)
TSHR_human	thyrotropin (TSHR) thyroid stimulating hormone	TMVII	677 CANPFLYAIFT SEQ ID NO: 58 V	cAMP formation / CHO cells	(Russo, Wong et al. 1999)
TSHR_human	thyrotropin (TSHR) thyroid stimulating hormone	IC3	613 VRNPQYNPGDKDTIAK deletion SEQ ID NO: 59 621	cAMP formation / COS-7	(Wonerow, Schoneberg et al. 1998)

FIG. 1-9

9/31

TSHR_human	thyrotropin (TSHR)	IC3 / TMVI	SEQ ID NO: 60	623 KDTKIAKRMVLIPTDFICM V I	cAMP activation / COS-7	(Paschke, Tonacchera et al. 1994)
V2R_human	thyroid stimulating hormone vasopressin V2	IC2	SEQ ID NO: 61	136 LMTLDRHRAI A	cAMP formation / COS-7	(Morin, Cotte et al. 1998)

10039645.050102

10/31

FIG. 1-10

File Name	Receptor	Mutation Site	Sequence	Assay / Cells	Reference
CLASS B GROUP I					
CALR_human	human calcitonin hCTR-1 hCTR-2	wild type (native) protein		adenylyl cyclase cAMP production / COS-1	(Cohen, Thaw et al. 1997)
CLASS B GROUP II					
PTRR_human	parathyroid hormone PTH / PTH-related peptide	junction of IC1 and TMII junction of IC3 and TMVI	223 TRNYIHHLFL SEQ ID NO: 62 R, K 410 KLLKSTLVLMF SEQ ID NO: 63 C, Others	cAMP accumulation / COS-7	(Schipani, Jensen et al. 1997)
CLASS B GROUP III					
GIPR_human	glucose-dependent insulinotropic peptide (GIP-R)	TMVI	340 VFAPVTEEQAR SEQ ID NO: 64 P	cAMP production / L293	(Tseng and Lin 1997)
GLR_rat	glucagon	junction of IC loop I and TMII IC end of TMVI	178 TRNYIHGNLFA SEQ ID NO: 65 R 352 RLARSTLTLP SEQ ID NO: 66 A	cAMP accumulation / COS-7	(Hjorth, Orskov et al. 1998)
VIPR_human	vasoactive intestinal peptide 1 (VIP)	junction of IC loop 1 and TMII junction of IC loop 3 and TMVI	178 RNYIHHLFI SEQ ID NO: 67 R functional integrity of the N-terminal EC domain 343 LARSTLTLP SEQ ID NO: 68 X= K, P	cAMP production / COS-7 or CHO	(Gaudin, Maoret et al. 1998) (Gaudin, Rouyer-Fessard et al. 1998)

FIG. 1-11

11/31

File Name	Receptor	Mutation Site	Sequence	Assay / Cells	Reference
CLASS C					
CASR_human	calcium-sensing	N-terminal EC	TLSFVAQNKIDSLNLDEECNCSEHA various substitutions, in multiple combinations SEQ ID NO: 69	IP / tsA	(Jensen, Spalding et al. 2000)

10029645 . 050102

FIG. 1-12

12/31

File Name	Receptor	Mutation Site	Sequence	Assay / Cells	Reference
CLASS D					
O74283 RCB2 <i>C. cinereus</i>	pheromone	TM6	229 PLSAYQIYLTGT SEQ ID NO: 70 P	heterologous yeast assay	(Olesnicky, Brown et al. 1999)
STE2_yeast	pheromone α -factor	TM6	258 QSLLVPSIIFI SEQ ID NO: 71 LL	<i>lacZ</i> reporter gene	(Konopka, Margarit et al. 1996)
STE2_yeast	pheromone α -factor	double mutations TM5 and TM6	223 MSFVLVVK Q ILAIR SEQ ID NO: 72 C C 247 251 DSFHILL Q SCSLL SEQ ID NO: 73 CC CC double mutations shaded double mutations	<i>lacZ</i> reporter gene / yeast	(Dube, DeCostanzo et al. 2000)
STE3_yeast	pheromone α -factor	IC3	194 DVRDILHCTNS SEQ ID NO: 74 Q	β -galactosidase	(Boone, Davis et al. 1993)
STE2_yeast	pheromone α -factor	TM6	253 258 LIMSCQSLLVPSIIFI SEQ ID NO: 75 L LP	β -galactosidase	(Sommers, Martin et al. 2000)

100W9645 . 05010N

FIG. 1-13

Bibliography

- Acharya, S. and S. S. Kamik (1996). "Modulation of GDP release from transducin by the conserved Glu134-Arg135 sequence in rhodopsin." J Biol Chem 271(41): 25406-11.
- Alewijnse, A. E., H. Timmerman, et al. (2000). "The Effect of Mutations in the DRY Motif on the Constitutive Activity and Structural Instability of the Histamine H(2) Receptor." Mol Pharmacol 57(5): 890-898.
- Allen, L. F., J. J. Lefkowitz, et al. (1991). "G-protein-coupled receptor genes as protooncogenes: constitutively activating mutation of the alpha 1B-adrenergic receptor enhances mitogenesis and tumorigenicity." Proc Natl Acad Sci U S A 88(24): 11354-8.
- Amaral, T. T., 3rd, S. Dragas-Graonic, et al. (1995). "Signal transduction by the formyl peptide receptor. Studies using chimeric receptors and site-directed mutagenesis define a novel domain for interaction with G-proteins." J Biol Chem 270(47): 28010-3.
- Hüml, K., E. Mutschler, et al. (1994). "Functional role in ligand binding and receptor activation of an asparagine residue present in the sixth transmembrane domain of all muscarinic acetylcholine receptors." J Biol Chem 269(29): 18870-6.
- Boone, C., N. G. Davis, et al. (1993). "Mutations that alter the third cytoplasmic loop of the α -factor receptor lead to a constitutive and hypersensitive phenotype." Proc Natl Acad Sci U S A 90(21): 9921-5.
- Bradbury, F. A., N. Kawate, et al. (1997). "Post-translational processing in the Golgi plays a critical role in the trafficking of the luteinizing hormone/human chorionic gonadotropin receptor to the cell surface." J Biol Chem 272(9): 5921-6.
- Bradbury, F. A. and K. M. Menon (1999). "Evidence that constitutively active luteinizing hormone/human chorionic gonadotropin receptors are rapidly internalized." Biochemistry 38(27): 8703-12.
- Burger, M., J. A. Burger, et al. (1999). "Point mutation causing constitutive signaling of CXCR2 leads to transforming activity similar to Kaposi's sarcoma herpesvirus-G protein-coupled receptor." J Immunol 163(4): 2017-22.
- Burstein, E. S., T. A. Spalding, et al. (1996). "Constitutive activation of chimeric m2/m5 muscarinic receptors and delineation of G-protein coupling selectivity domains." Biochem Pharmacol 51(4): 539-44.
- Cavalli, A., A. M. Babey, et al. (1999). "Altered adenylyl cyclase responsiveness subsequent to point mutations of Asp 128 in the third transmembrane domain of the delta-opioid receptor." Neuroscience 93(3): 1025-31.
- Charpentier, S., K. R. Jarvie, et al. (1996). "Silencing of the constitutive activity of the dopamine D1B receptor. Reciprocal mutations between D1 receptor subtypes delineate residues underlying activation properties." J Biol Chem 271(45): 28071-6.
- Cho, W., L. P. Taylor, et al. (1996). "Mutagenesis of residues adjacent to transmembrane prolines alters D1 dopamine receptor binding and signal transduction." Mol Pharmacol 50(5): 1338-45.
- Cohen, D. P., C. N. Thaw, et al. (1997). "Human calcitonin receptors exhibit agonist-independent (constitutive) signaling activity." Endocrinology 138(4): 1400-5.
- Cohen, G. B., T. Yang, et al. (1993). "Constitutive activation of opsin: influence of charge at position 134 and size at position 296." Biochemistry 32(23): 6111-5.
- Dube, P., A. DeCostanzo, et al. (2000). "Interaction between transmembrane domains five and six of the alpha -factor receptor." J Biol Chem 275(34): 26492-9.
- Duprez, L., J. Parma, et al. (1994). "Germline mutations in the thyrotropin receptor gene cause non-autoimmune autosomal dominant hyperthyroidism." Nat Genet 7(3): 396-401.
- Egan, C. T., K. Herrick-Davis, et al. (1998). "Creation of a constitutively activated state of the 5- hydroxytryptamine2A receptor by site-directed mutagenesis: inverse agonist activity of antipsychotic drugs." J Pharmacol Exp Ther 286(1): 85-90.
- Esapa, C. T., L. Duprez, et al. (1999). "A novel thyrotropin receptor mutation in an infant with severe thyrotoxicosis." Thyroid 9(10): 1005-10.
- Fanelli, F., P. Barbier, et al. (1999). "Activation mechanism of human oxytocin receptor: a combined study of experimental and computer-simulated mutagenesis." Mol Pharmacol 56(1): 214-25.
- Gaudin, P., J. J. Maoret, et al. (1998). "Constitutive activation of the human vasoactive intestinal peptide 1 receptor, a member of the new class II family of G protein-coupled receptors." J Biol Chem 273(9): 4990-6.
- Gaudin, P., C. Rouyer-Fessard, et al. (1998). "Constitutive activation of the human VIP1 receptor." Ann NY Acad Sci 865: 382-5.

FIG. 1-14

- Giovardhan, C. P. and D. D. Oprea (1994). "Active site-directed inactivation of constitutively active mutants of rhodopsin." J Biol Chem 269(9): 6524-7.
- Groblewski, T., B. Maigret, et al. (1997). "Mutation of Asn111 in the third transmembrane domain of the AT1A angiotensin II receptor induces its constitutive activation." J Biol Chem 272(3): 1822-6.
- Han, M., S. O. Smith, et al. (1998). "Constitutive activation of opsin by mutation of methionine 257 on transmembrane helix 6." Biochemistry 37(22): 8253-61.
- Hasegawa, H., M. Negishi, et al. (1996). "Two isoforms of the prostaglandin E receptor EP3 subtype different in agonist-independent constitutive activity." J Biol Chem 271(4): 1857-60.
- Herrick-Davis, K., C. Egan, et al. (1997). "Activating mutations of the serotonin 5-HT2C receptor." J Neurochem 69(3): 1138-44.
- Hjorth, S. A., C. Orskov, et al. (1998). "Constitutive activity of glucagon receptor mutants." Mol Endocrinol 12(1): 78-86.
- Högger, P., M. S. Shockey, et al. (1995). "Activating and inactivating mutations in N- and C-terminal 13 loop junctions of muscarinic acetylcholine Hm1 receptors." J Biol Chem 270(13): 7405-10.
- Hwa, J., R. Gavin, et al. (1997). "Synergism of constitutive activity in alpha 1-adrenergic receptor activation." Biochemistry 36(3): 633-9.
- Ishii, I., T. Izumi, et al. (1997). "Alanine exchanges of polar amino acids in the transmembrane domains of a platelet-activating factor receptor generate both constitutively active and inactive mutants." J Biol Chem 272(12): 7846-54.
- Jensen, A. A., T. A. Spalding, et al. (2000). "Functional importance of the Ala116-Pro136 region in the calcium-sensing receptor. CONSTITUTIVE ACTIVITY AND INVERSE AGONISM IN A FAMILY C G-PROTEIN-COUPLED RECEPTOR [In Process Citation]." J Biol Chem 275(38): 29547-55.
- Jin, J., G. F. Mao, et al. (1997). "Constitutive activity of human prostaglandin E receptor EP3 isoforms." British J Pharmacol 121: 317-23.
- Kjelsberg, M. A., S. Cotecchia, et al. (1992). "Constitutive activation of the alpha 1B-adrenergic receptor by all amino acid substitutions at a single site. Evidence for a region which constrains receptor activation." J Biol Chem 267(3): 1430-3.
- Konopka, J. B., S. M. Margalit, et al. (1996). "Mutation of Pro-258 in transmembrane domain 6 constitutively activates the G protein-coupled alpha-factor receptor." Proc Natl Acad Sci USA 93(13): 6764-9.
- Kusugi, S., C. Van Dop, et al. (1995). "Characterization of heterogeneous mutations causing constitutive activation of the luteinizing hormone receptor in familial male precocious puberty." Hum Mol Genet 4(2): 183-8.
- Kudo, M., Y. Osuga, et al. (1996). "Transmembrane regions V and VI of the human luteinizing hormone receptor are required for constitutive activation by a mutation in the third intracellular loop." J Biol Chem 271(37): 22470-8.
- Leikowitz, R. J., S. Cotecchia, et al. (1993). "Constitutive activity of receptors coupled to guanine nucleotide regulatory proteins." Trends Pharmacol Sci 14(8): 303-7.
- Liu, J., N. Blin, et al. (1996). "Molecular mechanisms involved in muscarinic acetylcholine receptor-mediated G protein activation studied by insertion mutagenesis." J Biol Chem 271(11): 6172-8.
- Marie, J., C. Koch, et al. (1999). "Constitutive activation of the human bradykinin B2 receptor induced by mutations in transmembrane helices III and VI." Mol Pharmacol 55(1): 92-101.
- Mason, D. A., J. D. Moore, et al. (1999). "A gain-of-function polymorphism in a G-protein coupling domain of the human beta 1-adrenergic receptor." J Biol Chem 274(18): 12670-4.
- Matus-Leibovich, N., D. R. Nussenzveig, et al. (1995). "Truncation of the thyrotropin-releasing hormone receptor carboxyl tail causes constitutive activity and leads to impaired responsiveness in Xenopus oocytes and A1T20 cells." J Biol Chem 270(3): 1041-7.
- Morin, D., N. Conte, et al. (1998). "The D136A mutation of the V2 vasopressin receptor induces a constitutive activity which permits discrimination between antagonists with partial agonist and inverse agonist activities." FEBS Lett 441(3): 470-5.
- Nanavicz, T., L. Wang, et al. (1996). "Thrombin receptor activating mutations. Alteration of an extracellular agonist recognition domain causes constitutive signaling." J Biol Chem 271(2): 702-6.
- Olesnicki, N. S., A. J. Brown, et al. (1999). "A constitutively active G-protein-coupled receptor causes mating self-compatibility in the mushroom *Coprinus*." Embo J 18(10): 2756-63.
- Parent, J. L., C. Le Gouill, et al. (1996). "Mutations of two adjacent amino acids generate inactive and constitutively active forms of the human platelet-activating factor receptor." J Biol Chem 271(14): 7949-55.

FIG. 1-15

- Parma, J., J. Van Sande, et al. (1995). "Somatic mutations causing constitutive activity of the thyrotropin receptor are the major cause of hyperfunctioning thyroid adenomas: identification of additional mutations activating both the cyclic adenosine 3',5'-monophosphate and inositol phosphate-Ca²⁺ cascades." Mol Endocrinol 9(6): 725-33.
- Parnot, C., S. Bardini, et al. (2000). "Systematic identification of mutations that constitutively activate the angiotensin II type 1A receptor by screening a randomly mutated cDNA library with an original pharmacological bioassay." Proc Natl Acad Sci U S A 97(13): 7615-20.
- Paschke, R., M. Tonacchera, et al. (1994). "Identification and functional characterization of two new somatic mutations causing constitutive activation of the thyrotropin receptor in hyperfunctioning autonomous adenomas of the thyroid." J Clin Endocrinol Metab 79(6): 1785-9.
- Pauwels, P. J., A. Gouble, et al. (1999). "Activation of constitutive 5-hydroxytryptamine 1B receptor by a series of mutations in the BBXXB motif: positioning of the third intracellular loop distal junction and its Gα protein interactions [In Process Citation]." Biochem J 343 Pt 2: 435-42.
- Perez, D. M., J. Hwa, et al. (1996). "Constitutive activation of a single effector pathway: evidence for multiple activation states of a G protein-coupled receptor." Mol Pharmacol 49(1): 112-22.
- Ren, Q., H. Kurose, et al. (1993). "Constitutively active mutants of the alpha 2-adrenergic receptor [published erratum appears in J Biol Chem 1994 Jan 14;269(2):1566]." J Biol Chem 268(22): 16483-7.
- Rim, J. and D. D. Orian (1995). "Constitutive activation of opsin: interaction of mutants with rhodopsin kinase and arrestin." Biochemistry 34(37): 11938-45.
- Robbins, L. S., J. H. Nadeau, et al. (1993). "Pigmentation phenotypes of variant extension locus alleles result from point mutations that alter MSH receptor function." Cell 72(6): 827-34.
- Russo, D., M. G. Wong, et al. (1999). "A Val 677 activating mutation of the thyrotropin receptor in a Hürthle cell thyroid carcinoma associated with thyrotoxicosis." Thyroid 9(1): 13-7.
- Samama, P., S. Cotecchia, et al. (1993). "A mutation-induced activated state of the beta 2-adrenergic receptor. Extending the ternary complex model." Journal of Biological Chemistry 268(7): 4625-36.
- Scheer, A., T. Costa, et al. (2000). "Mutational analysis of the highly conserved arginine within the Glu/Asp-Arg-Tyr motif of the alpha(1b)-adrenergic receptor: effects on receptor isomerization and activation." Mol Pharmacol 57(2): 219-31.
- Scheer, A., F. Fanelli, et al. (1997). "The activation process of the alpha 1B-adrenergic receptor: potential role of protonation and hydrophobicity of a highly conserved aspartate." Proc Natl Acad Sci U S A 94(3): 808-13.
- Schipani, E., G. S. Jensen, et al. (1997). "Constitutive activation of the cyclic adenosine 3',5'-monophosphate signaling pathway by parathyroid hormone (PTH)/PTH-related peptide receptors mutated at the two loci for Jansen's metaphyseal chondrodysplasia." Mol Endocrinol 11(7): 851-8.
- Shenker, A., L. Laue, et al. (1993). "A constitutively activating mutation of the luteinizing hormone receptor in familial male precocious puberty [see comments]." Nature 365(6447): 652-4.
- Soumiers, C. M., N. P. Martin, et al. (2000). "A limited spectrum of mutations causes constitutive activation of the yeast alpha-factor receptor." Biochemistry 39(23): 6898-909.
- Spalding, T. A., E. S. Bursstein, et al. (1998). "Identification of a ligand-dependent switch within a muscarinic receptor." J Biol Chem 273(34): 21563-8.
- Spalding, T. A., E. S. Bursstein, et al. (1997). "Constitutive activation of the m5 muscarinic receptor by a series of mutations at the extracellular end of transmembrane 6." Biochemistry 36(33): 10109-16.
- Tseng, C. C. and L. Lin (1997). "A point mutation in the glucose-dependent insulinotropic peptide receptor confers constitutive activity." Biochem Biophys Res Commun 232(1): 96-100.
- Wonerow, P., T. Schöneberg, et al. (1998). "Deletions in the third intracellular loop of the thyrotropin receptor. A new mechanism for constitutive activation." J Biol Chem 273(14): 7900-5.

FIG. 2
A Point Mutation Enhances MC-4 Receptor
Constitutive Activity




FIG. 3

Light Emission Induced by the WT CCK-BR
vs. a Constitutively Active Mutant





FIG. 4

A Point Mutation Confers Constitutive Activity to the Rat μ Opioid Receptor



19/31

FIG. 5

Forskolin Stimulated HEK293 Cells Transfected
With pcDNA1 and a CRE-luc Construct




FIG. 6

The Rat μ Opioid Receptor Signals Through G α i





FIG. 7

A Point Mutation Confers Constitutive Activity to the Rat μ Opioid Receptor



Target Residues Within Class I GPCRs





FIG. 9


TMD III Asn (-14 from DRY) is a Target
for Mutation Induced Constitutive Activity



24/31

FIG. 10

The 'DRY' Motif is a Target for Mutation
Induced Constitutive Activity



25/31

FIG. 11

A Point Mutation Enhances MC-4 Receptor
Constitutive Activity





FIG. 12

The -13 Position is a Target for Mutation
Induced Constitutive Activity



27/31

FIG. 13

SEQ ID NO: 76 ork 1 -----MESPIQIFRGEPEGTCAPSACLPNSSAWFPGWAF..DSNGSAGSEDAG
 SEQ ID NO: 77 orkr 1 -----MESPIQIFRGEPEGTCAPSACLPNSSSWFPNWAES..DSNGSVGSEDOO
 SEQ ID NO: 78 orm 1 MDSSAAPTNASNCTDAAYSSCSAPSEGSWV..NLSHLDGMLSDPCGNRTDLGGRLSL
 SEQ ID NO: 79 ormr 1 MDSSTGPGNTSDCSDPIAQASCSEA..PGSWL..NLSHVDGNQSDPCGLNRTGLGNDLSL
 SEQ ID NO: 80 ord 1 -----MEAPASAGAE..PPLFANASDAYPSACPSACANASG
 SEQ ID NO: 81 ATla 1 -----MALNSSAEDGIKRIG
 SEQ ID NO: 82 BK-2 1 -----MFSPWKISMFLSVREDSVPTTASFSADMLNVTLQGP TLNG.TFAC

ork 49 LEPANISPAH..PVHITAYYSVVFVVGLAGNSLVMVITRYTKMKTATNIYIFNLALADA
 orkr 49 LEPANISPAH..PVHITAYYSVVFVVGLAGNSLVMVITRYTKMKTATNIYIFNLALADA
 orm 59 CPPTGS.PSMITAITIMALYSIVCVVGLFGNFLVMVIMRYTKMKTATNIYIFNLALADA
 ormr 57 CPQTGS.PSMVTAITIMALYSIVCVVGLFGNFLVMVIMRYTKMKTATNIYIFNLALADA
 ord 37 PPGASASSIALATAITIMALYSIVCAVGLFGNFLVMVIMRYTKMKTATNIYIFNLALADA
 ATla 16 DDCPMAGRHSYIFVMIPTLYSICFVVGIFCNLSLVIVVIFYFMKKTIVASVFLNLALADL
 BK-2 45 SKCPQVEWLGLNLTIPPPFLWVEFLATENIFVLSVFLCHKSSCIVAETVLCNLAAADL

ork 107 LVTHITMPFQSTVYLMN..SWPFGDVLCKIVISIDYNNMFTSIFTLTMSVDRIYAVCHPVK
 orkr 107 LVTHITMPFQSAVYLMN..SWPFGDVLCKIVISIDYNNMFTSIFTLTMSVDRIYAVCHPVK
 orm 118 LATSTLPPFQSVNYLNG..TWPFGLTLCKIVISIDYNNMFTSIFTLTMSVDRIYAVCHPVK
 ormr 116 LATSTLPPFQSVNYLNG..TWPFGLTLCKIVISIDYNNMFTSIFTLTMSVDRIYAVCHPVK
 ord 97 LATSTLPPFQSAKYLMZ..TWPFGEHLCKAVESIDYNNMFTSIFTLTMSVDRIYAVCHPVK
 ATla 76 CFLTLPLWAVYTAMEYRWPFGNHLCKIASAVTENLYASVLTCTSDRYIAIVHPMK
 BK-2 105 ILACGLPFWAITISNNFDWLGETLCRVNMTISMNLYSSICFMDVMSIDRYIAIVKIMS

-14 from DRY *

ork 166 ALDFRTEFLKAKIINICIWLLSSVGSISAVLGGTKVR..EDVDVIECSLOFEDDDYSND
 orkr 166 ALDFRTEFLKAKIINICIWLLSSVGSISAVLGGTKVR..EDVDVIECSLOFEDDEYSND
 orm 177 ALDFRTERNAKINVCNWILSSAIGLPVFMATTIKVR..Q..GSIDCTLTSHTPTW..KVE
 ormr 175 ALDFRTERNAKINVCNWILSSAIGLPVFMATTIKVR..Q..GSIDCTLTSHTPTW..KVE
 ord 156 ALDFRTEAKAKIINICIWVLSAIGVGPVIMVAVTPR..D..GAVVOMLOFESPSW..YWD
 ATla 136 SRLRRIMLVAKVTCIITWLVAGLASPAVHRNV..YFIENNTMTVCAFRYESRN.STLP
 BK-2 165 MGRMRGVRWAKLYSIVWGCALLSSPMLVFRMTKEYSDEGHNVACVHSYS...LIVE

ork 224 IFMKICVFIFAFVTPVLIITVCYTLMLRLKSVRLSGSREKDRNLRRITRLVLVVAVF
 orkr 224 IFMKICVFIFAFVTPVLIITVCYTLMLRLKSVRLSGSREKDRNLRRITRLVLVVAVF
 orm 232 NLKICVFIFAFVTPVLIITVCYGLMLRLKSVRLSGSKEKDRNLRRITRMVLVVAVF
 ormr 230 NLKICVFIFAFVTPVLIITVCYGLMLRLKSVRLSGSKEKDRNLRRITRMVLVVAVF
 ord 211 TVTKICVFIFAFVTPVLIITVCYGLMLRLKSVRLSGSKEKDRSLRRITRMVLVVAVF
 ATla 193 TGLGCTKNILGFLFPLIITLSYTLIWKALKKAYEIOKNKPRND...IFRIIMATVLF
 BK-2 222 VFTNMLNIVVGFLHP..LSVITFCITMOIMVLRNNEYOKFKEIQTE..RRATVVLVLVLLHF

ork 284 IVCWTPIHIFILVHALGS.T.....SHSTAALSSYFCIALGYTNSSLNPLVYAFLDENF
 orkr 284 IVCWTPIHIFILVHALGS.T.....SHSTAALSSYFCIALGYTNSSLNPLVYAFLDENF
 orm 292 IVCWTPIHIYVILKALVTIP.....ETTFQTVSWHFCIALGYTNSSLNPLVYAFLDENF
 ormr 290 IVCWTPIHIYVILKALVTIP.....ETTFQTVSWHFCIALGYTNSSLNPLVYAFLDENF
 ord 271 IVCWTPIHIFIVITLVLDID.....RRDPLVVAALHFCIALGYANSSLNPLVYAFLDENF
 ATla 250 FFSWVPHQIETFLDVLITOLGVHDKISDIVDTAMPITICTAYENNCNPLIYGFGLGKKE
 BK-2 280 IICWLPFQISTFIDTLHRLGILSSCODERIIDVITQIASPMAYSNSCLNPLVYVIVGKRE

ork 338 KRCFRDFCFPLKMRMEROSTSRVR.NTVOD..PAYLRDIEDGMNKPV-----
 orkr 338 KRCFRDFCFPLKMRMEROSTSRVR.NTVOD..PASMEDVGGMKNKPV-----
 orm 346 KRCFRDFCFIPTSSNIEQONSTRFRONT..RDHPSANTVDRTNHOLENLEAETAPLP
 ormr 344 KRCFRDFCFIPTSSNIEQONSTRFRONT..RDHPSANTVDRTNHOLENLEAETAPLP
 ord 326 KRCFRDORCPGCPDPSFSHAREATAREVRTACTPSDGPGGGAAA-----
 ATla 310 KKMFLQLLKYPKAKSHS...SLSTKM..STLSYRPSDNSSAKKPASCFEVE-
 BK-2 340 RKKSWEVYQGVCGGGCRSEPIQMENS..GTL..RTSISVDRTHKLQDWAGSRO

FIG. 14

28/31

SEQ ID NO: 83 mOrmouse 1 MDSSAGPGNITSDCSPLA.PASCSPA..PGSWNLHSHVDGNOSDPCGPNRTGLGGSHSLG
 SEQ ID NO: 79 mOrRat 1 MDSSITGPGNITSDCSPLA.OASCSPA..PGSWNLHSHVDGNOSDPCGPNRTGLGGNDLSLQ
 SEQ ID NO: 84 mOrbovin 1 MDSCAVPTNANCTDDEFFTHPSSCSPAPSPSSWVNFSLHLCNLSDPCGPNRTGLGGSDRLC
 SEQ ID NO: 85 mOrhuman 1 MDSSAAPTNANCTDDEFFTHPSSCSPAPSPSSWVNFSLHLCNLSDPCGPNRTGLGGSDRLC
 SEQ ID NO: 86 mOrpig 1 MDSSADPRNANCTDDEFFSPSSMCSPVPSPSSWVNFSLHLCNLSDPCGPNRTGLGGSDSLC
 SEQ ID NO: 87 mOrws 1 METS...GNISDFLYPLS.....NEVMS.....NSSVLCRNFSNSTSFLNMGSSRSDTD
 SEQ ID NO: 81 AT1a 1 -----MALNSSAEDGKRIQDDC
 SEQ ID NO: 82 BK-2 1 -----MFSFWKISMFLSVREDSVPTTASFSAFLNVTLOGETLNG.TFAQSKG

mOrmouse 58 POTGSPSMUTAITIMALYSIVCVVGLFGNFLVMYVIVRYTKMKATATNIYIFNLALADALA
 mOrRat 58 POTGSPSMUTAITIMALYSIVCVVGLFGNFLVMYVIVRYTKMKATATNIYIFNLALADALA
 mOrbovin 61 PSAGSPSMUTAITIMALYSIVCVVGLFGNFLVMYVIVRYTKMKATATNIYIFNLALADALA
 mOrhuman 60 PPTGSPSMUTAITIMALYSIVCVVGLFGNFLVMYVIVRYTKMKATATNIYIFNLALADALA
 mOrpig 61 PPTGSPSMUTAITIMALYSIVCVVGLFGNFLVMYVIVRYTKMKATATNIYIFNLALADALA
 mOrws 48 EODKLP.VITAITITILYSIVCVVGLVGNVLMYVIVRYTKMKATATNIYIFNLALADALA
 AT1a 19 PKAGRHSYIFVM.IPTLYSIHFVVGIFGNSLVVIVRYTKMKATATNIYIFNLALADALA
 BK-2 48 POVEWLGWNTI.QPPFLWVIFVLETLNIFVLSVFLCHKSSCTVAETIYIENLAADLIL

mOrmouse 118 TSTLPFQSVNYLMG.TWPFGNILCKIVISIDYNNMETSIFLTCTMSVDRIYAVCHPVKAL
 mOrRat 118 TSTLPFQSVNYLMG.TWPFGNILCKIVISIDYNNMETSIFLTCTMSVDRIYAVCHPVKAL
 mOrbovin 121 TSTLPFQSVNYLMG.TWPFGNILCKIVISIDYNNMETSIFLTCTMSVDRIYAVCHPVKAL
 mOrhuman 120 TSTLPFQSVNYLMG.TWPFGNILCKIVISIDYNNMETSIFLTCTMSVDRIYAVCHPVKAL
 mOrpig 121 TSTLPFQSVNYLMG.TWPFGNILCKIVISIDYNNMETSIFLTCTMSVDRIYAVCHPVKAL
 mOrws 107 TSTLPFQSVNYLMG.TWPFGNILCKIVISIDYNNMETSIFLTCTMSVDRIYAVCHPVKAL
 AT1a 78 LLTLEPLWAVYTAMEYRWPFGNILCKIVISIDYNNMETSIFLTCTMSVDRIYAVCHPVKAL
 BK-2 107 ACGLPEWATITISNNFDMFGETLCEVVMIIISMNLYSSICFLMLSSIDRIYAVCHPVKAL

mOrmouse 177 DFRTPRNAKIVNVCNWILSSAIGLPVMFMATTKYRC.....GSIDCTLTFSHPTWYWE
 mOrRat 177 DFRTPRNAKIVNVCNWILSSAIGLPVMFMATTKYRC.....GSIDCTLTFSHPTWYWE
 mOrbovin 180 DFRTPRNAKIVNVCNWILSSAIGLPVMFMATTKYRC.....GSIDCTLTFSHPTWYWE
 mOrhuman 179 DFRTPRNAKIVNVCNWILSSAIGLPVMFMATTKYRC.....GSIDCTLTFSHPTWYWE
 mOrpig 180 DFRTPRNAKIVNVCNWILSSAIGLPVMFMATTKYRC.....GSIDCTLTFSHPTWYWE
 mOrws 166 DFRTPRNAKIVNVCNWILSSAIGLPVMFMATTKYRC.....GSIDCTLTFSHPTWYWE
 AT1a 138 LRRIMLVAKVTCIIIMMAGLASLBAVIHRNV....YFIENNTITVAFHYESRNSTLP
 BK-2 167 RMRGVVWAKVYSLVINGCILLSSPMLVFRIMF...EYSDEGHNVTAQVLSYPS..LINE

mOrmouse 230 NLLKICVFIFAFIMPVLIITVCYGLMILRLKSVRMLSGSKEKDRNLRRITRMVLVVAVF
 mOrRat 230 NLLKICVFIFAFIMPVLIITVCYGLMILRLKSVRMLSGSKEKDRNLRRITRMVLVVAVF
 mOrbovin 233 NLLKICVFIFAFIMPVLIITVCYGLMILRLKSVRMLSGSKEKDRNLRRITRMVLVVAVF
 mOrhuman 232 NLLKICVFIFAFIMPVLIITVCYGLMILRLKSVRMLSGSKEKDRNLRRITRMVLVVAVF
 mOrpig 233 NLLKICVFIFAFIMPVLIITVCYGLMILRLKSVRMLSGSKEKDRNLRRITRMVLVVAVF
 mOrws 226 TLLKICVFIFAFIMPVLIITVCYGLMILRLKSVRMLSGSKEKDRNLRRITRMVLVVAVF
 AT1a 193 IGLGETKNILGFDFPFLIILTSYTLIWKALKAYETOKNKPND...LRTIMAVLFF
 BK-2 222 VFTNMLNVLVGLLP.LSVITFCTYQIMVLRNNEQKFKETOTE.RRATVLVLVLLFF

mOrmouse 290 IVCWTPIHIYVIAKALITI.....PETTFQTVSWHFCIALGYTNSCLNPVLYAFLDENF
 mOrRat 290 IVCWTPIHIYVIAKALITI.....PETTFQTVSWHFCIALGYTNSCLNPVLYAFLDENF
 mOrbovin 293 IVCWTPIHIYVIAKALITI.....PETTFQTVSWHFCIALGYTNSCLNPVLYAFLDENF
 mOrhuman 292 IVCWTPIHIYVIAKALITI.....PETTFQTVSWHFCIALGYTNSCLNPVLYAFLDENF
 mOrpig 293 IVCWTPIHIYVIAKALITI.....PETTFQTVSWHFCIALGYTNSCLNPVLYAFLDENF
 mOrws 286 IVCWTPIHIYVIAKALITI.....PETTFQTVSWHFCIALGYTNSCLNPVLYAFLDENF
 AT1a 250 FFSWVPHQISTFDVLIQGVIIHDKISDIVDTAMPITICTAYENNCLNPLFYGFLGKKF
 BK-2 280 IICRLIFQISTFDTHRGILSSCODERIIDVITQIASFMVNSCLNPVLYVIGKRF

mOrmouse 344 KRCFREFC..IPTSSSTIEQONSARIRONTRHPSTANTVDRTNHOLENLEAETAPLF
 mOrRat 344 KRCFREFC..IPTSSSTIEQONSARIRONTRHPSTANTVDRTNHOLENLEAETAPLF
 mOrbovin 347 KRCFREFC..IPTSSSTIEQONSARIRONTRHPSTANTVDRTNHOLENLEAETAPLF
 mOrhuman 346 KRCFREFC..IPTSSSTIEQONSARIRONTRHPSTANTVDRTNHOLENLEAETAPLF
 mOrpig 347 KRCFREFC..IPTSSSTIEQONSARIRONTRHPSTANTVDRTNHOLENLEAETAPLF
 mOrws 340 KRCFREFC..VPSFVLDONSTRNSNPOCGCGSGHKVDRNTRGV-----
 AT1a 310 KKYFLCLLKYIPPAKSHS...SLSTKSTLSYRPSDMSSSAKPAFCFEVE----
 BK-2 340 RKKSWEVYQGVQKGGCRSEPIOMENSMGTL...RTSISVERQIKLQDWAGSRO----

29/31

FIG. 15





FIG. 16

An Intracellular Point Mutation Results in
Loss of Ligand-Induced Function



31/31

FIG. 17

